

Boosters Final CDC /FDA approval received

Note: boosters may be given of a different vaccine than originally given, dosing schedule will be according to the original vaccine given. For example, If original vaccine was J and J, booster with Moderna can be given at 2 months after original dose of J and J. If original vaccine was Moderna and they want J and J, it would be given 6 months after the second Moderna vaccine.

Product	Amount	Timing	Who can receive
Pfizer-Comernity	Same as original dose	6 months after 2 nd dose	65 and older, 18 + resident and emps of LTC, healthcare workers, workers in high risk environments, 18+ with underlying medical conditions
Moderna	½ original dose	6 months after 2 nd dose	65 and older, 18 + resident and emps of LTC, healthcare workers, workers in high risk environments, 18+ with underlying medical conditions
J and J	Same as original dose	2 months or longer after 1 st dose	Anyone that has received the first dose

3rd dose (NOT A BOOSTER) of Pfizer or Moderna is for immunocompromised persons to be given 28 days after second dose:

These are the conditions for its use:

You may give the third dose of vaccine to patients or staff that meet the following criteria:

- Receiving active cancer treatment for tumors or cancers of the blood;
- Received an organ transplant and am taking medicine to suppress my immune system;
- Received a stem cell transplant within the last two years or am taking medicine to suppress the immune system;
- Moderate or severe primary immunodeficiency (such as DiGeorge syndrome, Wiskott-Aldrich syndrome);
- Advanced or untreated HIV infection;
- Active treatment with high-dose corticosteroids or other drugs that suppress my immune response